

Profiler Applications

Wayne Angevine

CIRES / NOAA Aeronomy Lab

I. Boundary layer height

II. Temperature (RASS)

Allen White

CIRES / NOAA Environmental Technology Lab

III. Turbulence intensity

IV. Weather phenomena

Profiler Applications


I. Boundary layer height

- Orientation
- Reflectivity patterns
- Complications
- More information to aid interpretation

II. Temperature (RASS)

- Accuracy, precision, and biases
- Applications

Atmospheric Boundary Layer Diurnal Variation




Adapted from *Introduction to Boundary Layer Meteorology* -R.B. Stull, 1988

How is BL height found?

- Primary method uses reflectivity pattern
 - Reflectivity peaks at BL top, providing a sharp indication
- Other parameters are used to clarify difficult or ambiguous cases
 - Ceilometer cloud base
 - Spectral width
 - RASS temperature

How does a profiler see the ABL?


- Reflectivity is roughly the product of humidity gradient and turbulence intensity



Deep layer

- Note deep entrainment signatures



Day 187 1996




Residual layer

- Can fool simple automated algorithms

Day 253 1995





Reflectivity, PBL height, and cloud base



Low humidity

➤ Weakens signal


Day 266 1995



Weak inversion, cloud

- Broad or weak inversion
- “Deep” but non-precipitating cloud
- Ceilometer aids greatly in interpretation


Day 228 1995



Weak inversion, cloud

- Note disappearance of BL signature 1400-1500 CST



Day 207 1996



Multiple layers

- Note correlation of layers to soundings



Day 218 1996




Multiple layers

- Can fool simple automated algorithms

Day 233 1995





Reflectivity, PBL height, and cloud base





Overcast and rain

➤ (Pease day 196 2002)





Coastal BL with sea breeze

➤ (Pease day 215 2002)





Marine BL

➤ (Appledore day 181 2002)



Marine BL

- With advected continental BL in the morning
- (Appledore day 215 2002)



Basic rule

- Profilers can find convective boundary layer when it is defined physically
- Other types of BLs are more challenging
- Sometimes the physical definition of the BL is the issue

How do we find BL height in practice?

- Most reliable method is subjective determination based on reflectivity pattern with all available ancillary measurements
- Some automated algorithms exist in research environments
- Using more information (cloud data, more moments, time & space patterns) yields better results


Morning transition

- Transition between nocturnal, stable BL and daytime convective BL
- Profiler reflectivity patterns give a very good indication of timing of morning transition

Afternoon transition



- Transition between fully-developed daytime convective ABL and nocturnal ABL
- Timing and shape of transition are critical to initiation of inertial oscillation / low-level jet, nighttime transport, distribution of pollutants, etc.
- Turbulence gradually decreases in height and intensity
- Reflectivity can be misleading, it tends to show the residual inversion
- Spectral width is most useful to distinguish active turbulent region from developing residual layer

Profiler reflectivity and spectral width patterns for a “typical” day









Spatial variation of BL height

- Urban dome or urban heat island measured by profilers in urban core and in surrounding rural areas
- Implications for pollutant concentration and transport




Coastal influence on BL behavior

- Five profilers in Houston area for TEXAQS2000
- LaMarque, near Galveston Bay, shows coastal influence on this day
- We are studying sea breeze influence on pollution episodes
- Also note non-ICRA data at Ellington



RASS

- Many profilers have RASS (Radio Acoustic Sounding System) attached to measure (virtual) temperature
- RASS data can be useful but their limitations must be understood
- Height coverage is limited and varies with wind speed and other conditions
- Height-dependent bias is present due to range error – correction algorithms exist but are non-trivial
- RASS combined with surface data can be useful for finding nocturnal BL heights



References

- Angevine, W.M., A.B. White, and S.K. Avery, 1994: Boundary layer depth and entrainment zone characterization with a boundary layer profiler. *Boundary Layer Meteor.*, **68**, 375-385.
- Angevine, W.M., W.L. Ecklund, D.A. Carter, K.S. Gage, and K.P. Moran, 1994: Improved radio-acoustic sounding techniques. *J. Atmos. Oceanic Technol.*, **11**, 42-49.
- Angevine, W.M., and W.L. Ecklund, 1994: Errors in radio acoustic sounding of temperature. *J. Atmos. Oceanic Technol.*, **11**, 837-848.
- Angevine, W.M., and J.I. MacPherson, 1995: Comparison of wind profiler and aircraft wind measurements at Chebogue Point, Nova Scotia. *J. Atmos. Oceanic Technol.*, **12**, 421-426.
- Carter, D.A., K.S. Gage, W.L. Ecklund, W.M. Angevine, P.E. Johnston, A.C. Riddle, J. Wilson, and C.R. Williams, 1995: Developments in UHF lower tropospheric wind profiling at NOAA's Aeronomy Laboratory. *Radio Sci.*, **30**, 977-1001.
- Peters, G., and W.M. Angevine, 1996: On the correction of RASS-temperature errors due to turbulence. *Contributions to Atmospheric Physics*, **69**, 81-96.
- Riddle, A.C., W.M. Angevine, W.L. Ecklund, E.R. Miller, D.B. Parsons, D.A. Carter, and K.S. Gage, 1996: In situ and remotely sensed horizontal winds and temperature intercomparisons obtained using Integrated Sounding Systems during TOGA COARE. *Contributions to Atmospheric Physics*, **69**, 49-62.

References, continued

- Angevine, W.M., 1997: Errors in mean vertical velocities measured by boundary layer wind profilers. *J. Atmos. Oceanic. Technol.*, **14**, 565-569.
- Angevine, W.M., P.S. Bakwin, and K.J. Davis, 1998: Wind profiler and RASS measurements compared with measurements from a 450 m tall tower. *J. Atmos. Oceanic. Technol.*, **15**, 818-825.
- Angevine, W.M., A.W. Grimsdell, L.M. Hartten, and A.C. Delany, 1998: The Flatland boundary layer experiments. *Bull. Amer. Meteorol. Soc.*, **79**, 419-431.
- Grimsdell, A.W., and W.M. Angevine, 1998: Convective boundary layer height measured with wind profilers and compared to cloud base. *J. Atmos. Oceanic Technol.*, **15**, 1332-1339.
- Görsdorf, U., and V. Lehmann, 2000: Enhanced accuracy of RASS-measured temperatures due to an improved range correction. *J. Atmos. Oceanic Technol.*, **17**, 406-416.
- Cohn, S.A., and W.M. Angevine, 2000: Boundary layer height and entrainment zone thickness measured by lidars and wind profiling radars. *J. Appl. Meteorol.*, **39**, 1233-1247.
- Angevine, W.M., H. Klein Baltink, and F.C. Bosveld, 2001: Observations of the morning transition of the convective boundary layer. *Boundary-Layer Meteorol.*, **101**, 209-227.
- Grimsdell, A.W., and W.M. Angevine, 2002: Observations of the afternoon transition of the convective boundary layer. *J. Appl. Meteorol.*, **41**, 3-11.
- Cohn, S.A., and R.K. Goodrich, 2002: Radar wind profiler radial velocity: A comparison with Doppler lidar. *J. Appl. Meteorol.*, **41**, 1277-1282.